

GLOBAL ESTIMATES OF OCCUPATIONAL ACCIDENTS AND WORK-RELATED ILLNESSES 2017

Päivi Hämäläinen . Jukka Takala . Tan Boon Kiat



Global Estimates of Occupational Accidents and Work-related Illnesses 2017

Published in September 2017 by
Workplace Safety and Health Institute
1500 Bendemeer Road, #04-01
Ministry of Manpower Services Centre
Singapore 339946
www.wsh-institute.sg

Copyright © 2017 Workplace Safety and Health Institute

RPT-2-2017

ISBN: 9789811148446

All rights reserved. This publication may not be reproduced or transmitted in any form or by any means, in whole or in part, without prior written permission.

CONTENT

Executive Summary	4
1 Background	5
2 Methodology	6
2.1 General.....	6
2.2 Fatal Occupational Accidents.....	8
2.3 Non-fatal Occupational Accidents.....	9
2.4 Fatal Work-Related Diseases.....	9
3 Results	11
3.1 Overview	11
3.2 Occupational Accidents	12
3.2.1 By WHO Division	12
3.2.2 By Geographical Region	12
3.3 Work-Related Diseases	13
3.3.1 Work-related Mortality by Cause	13
3.3.2 Number of Work-related Mortality by United Nations Geographical Regions	15
3.3.3 Deaths attributed to hazardous substances including dusts, vapours and fumes	16
4 Discussion	17
References	19

EXECUTIVE SUMMARY

This report provides an update to the global estimates of occupational accidents and work-related diseases that were presented during the XX World Congress on Safety and Health at Work 2014 at Frankfurt. These updated figures were released during the XXI World Congress held in Singapore on 3-6 Sep 2017, as agreed under the Memorandum of Understanding between the International Labour Organization (ILO) and the Workplace Safety and Health (WSH) Institute, Singapore. These estimates were worked out by a team comprising experts from the Ministry of Health and Social Affairs in Finland and Workplace Safety and Health Institute of the Ministry of Manpower in Singapore.

The methodology used in this update was the same as for the 2014 study so as to keep the estimated numbers comparable. In the 2014 study, estimates were made based on the World Health Organization (WHO) regions. In addition to the WHO classification, estimates were also made based on the United Nations (UN) geographical regions. As there was under-estimation on the number of respiratory disease cases in 2014, new attributable fractions (AF) were used to estimate the number of respiratory cases caused by Chronic Obstructive Pulmonary Disease (COPD) and Asthma.

We estimated 2.78 million deaths occurring annually across the countries being attributed to work, higher than the 2.33 million deaths estimated in 2014. Work-related mortality accounted for 5% of the global total deaths (based on the Global Burden of Disease Study 2015). The biggest share of work-related mortality came from work-related diseases which accounted for 2.4 million (86.3%) of the total estimated deaths. Fatal accidents accounted for the remaining 13.7%.

With the inclusion of COPD in our estimation, respiratory diseases (17%) had increased and was among the top three illnesses after circulatory diseases (31%) and malignant neoplasms (26%). Together, they contributed more than three-quarter of the total work-related mortality, followed by occupational injuries at 14% and communicable diseases (9%).

Asia was the highest contributor and constituted about two-thirds of the global work related mortality, followed by Africa at 11.8% and Europe at 11.7%.

Similar to previous estimates, the number of estimated work-related illnesses far exceeds that of work accident and fatalities. The increase in the number of ill-health was due to previous underestimates of COPD, leading to the respiratory estimates climbing to the third spot among the work related mortality. Asia emerges top and constitutes about two-thirds of the estimated work-related mortality, nearly 6 times that of Africa and Europe.

1 Background

This report is an update to the global estimates of occupational accidents and work-related diseases that was shared during the XX World Congress at Frankfurt in 2014. A Memorandum of Understanding (MOU) was signed between the International Labour Organization (ILO) and Workplace Safety and Health (WSH) Institute, Singapore to produce and release these estimates during the XXI World Congress in Sep 2017.

Tampere University of Technology (TUT) had been collaborating with ILO for over a decade to produce global estimates of fatal and non-fatal occupational accidents for 1998, 2001, 2003, 2008 and 2010

^{1,2} Global estimates of the fatal work-related diseases were also produced for the years 2000, 2002, 2008 and 2011. The latest scientific peer-reviewed report (Takala et al. 2014) was published by the Journal of Occupational and Environmental Hygiene in May 2014. For this latest report, the WSH Institute collaborated with the Finnish Ministry of Social Affairs and Health, OSHD Finland to produce the estimates for ILO.

For this project, estimates of the occupational accidents were compiled using 2014 data while the work-related diseases were based on 2015 data. Besides presenting the findings based on the seven World Health Organization (WHO) regions, findings were also presented based on the geographical and sub-regional groupings of the United Nations (UN). As the number of respiratory diseases had been underestimated previously, new attributable fractions (AF) were thus used in the estimation of the work-related diseases caused by Chronic Obstructive Pulmonary Disease (COPD) and Asthma.

2 Methodology

2.1 General

Sources of data on the burden of injuries and illnesses at work

We reviewed employment figures, mortality rates, occupational burden of diseases and injuries, reported accidents, surveys on self-reported occupational illnesses and injuries, attributable fractions, and the most recent information on the problems from published papers, documents, and electronic data sources of international and regional organizations, in particular ILO, WHO, the European Union (EU) and the Association of Southeast Asian Nations (ASEAN), institutions, agencies, and public websites.

Occupational injuries and work-related illnesses

While it is difficult to compare national data related to occupational injuries due to differences in legal and compensation criteria, the comparison between the number of fatal injuries (accidents) is easier and, although not completely, it is relatively comparable when the recording criteria, denominators and economic structures are well documented. Usually fatal injuries are expressed per 100,000 employed population in national statistics, or per 1 million working hours which may be converted to 100,000 full time employed as indicated in the Chinese National statistics.³

Occupational disease is usually understood to be a disorder or illness included in the designated list of diseases at work where employment injury compensation is legally defined. Inclusion in such a list is influenced by both evidence of work-relatedness and by administrative, financial and political arguments by the government or region concerned. In reality, the number of illnesses covered depends on the countries' development, political commitment and financial capacity. A disease, such as musculoskeletal disorder may be included in a list of occupational diseases in a country while not included in the list for another country. The concept of occupational injuries is usually much better defined, compared to occupational diseases.

Work-relatedness is a gradual component that may vary between obvious and commonly agreed to barely detectable. Often the work-relatedness is measured by the population attributable fraction (AF).

Past studies on work-related illnesses, and often including injuries, have been carried out in several countries, and globally. These have been based on the attributable fraction method and cover a varying number of illnesses and diseases. The AF method is well described by e.g. Nurminen and Karjalainen⁴. Detailed information on the attributable fractions and general methodology are available in Hämäläinen^{5,6}. This method has been used and applied including the AF values in Spain⁷ and New Zealand⁸ and with varied AF values in the United States⁹ and United Kingdom¹⁰. In an earlier comparison of different estimation methods the method used in this study was confirmed as the most realistic one.¹¹

Methods and data sources for the estimates related to fatal outcomes

Methods used in collecting data for the ILO, WHO and Global Burden of Disease, (GBD) of the Institute of Health Metrics and Evaluation (IHME) are explained elsewhere^{12,13,14,15,16}. The denominator in fatal injury rates is usually the number of employed persons. However, in other cases, the full labour force or data on the total population by age groups had been used. In many developing countries, the number of those formally employed is a small fraction of all who work in the country as self-employed, subsistence farmers, and those working in the informal sectors were usually not captured in the statistics. Globally, the labour force size may also be an underestimate in some countries as large numbers of those below 15 years of age, starting from as young as under ten years old and a sizeable number of those beyond any retirement age must work for survival. The same applies to collecting data on injury incidents.

Table 1. Summary of ILO method and data sources¹⁷

Estimates of work-related deaths	Methods/ Data Sources
a) Total number of deaths due to work injuries	<p>No. of fatal injuries reported to the ILO and member States</p> <ul style="list-style-type: none"> - Included fatal injuries, injury rates, work-related traffic fatalities suicides, but excluded fatalities via commuting to work and back - As ILO data includes data from a limited number of countries, those countries where no information was available were grouped in specific regions, in particular, WHO regions and sub-regions, and fatal injury rates per 100,000 employed of one or several countries that had produced injury rates were used as proxy values, - To increase the accuracy, separate injury rates were used for 1) agriculture and fishing, 2) industry including construction, and 3) service sector. This balances some of the potential differences between reporting proxy countries and non-reporting countries.
b) Total number of deaths due to work-related diseases	<p>All-cause mortality data from the World Health Organization</p> <ul style="list-style-type: none"> - Breakdown by diseases and groups of diseases - Breakdown by gender and age groups <p>Applied disease-specific AFs to the all-cause mortality data in 2015.</p> <p>The AFs were largely referenced from the Nurminen 2001 paper, with some replaced by more recent AFs. Related epidemiological studies and for selected industries and occupations to which the AFs were derived, were also reviewed. When no systematic data existed from elsewhere on the number and share of exposed workers, figures for similar occupations from Nurminen were used.</p> <p>Certain conditions were applied to exclude non-work-related deaths. For instance, children were excluded by using only deaths between the specific age groups</p>
c) Total number of suicides due to work	<p>Applied AF in Nurminen 2001 data to mortality data from WHO</p>

2.2 Fatal Occupational Accidents

The number of fatal occupational accidents was estimated from the 2014 frequency rates of fatal accidents (fatalities per 100 000 workers) from selected ILO member States that reported their accident data in three economic sectors:

- Agriculture including farming, fishing and forestry
- Industry including mining, manufacturing, energy production and construction
- Services

For countries where fatal data was not available, the WHO regions were used, where countries of similar income structures were grouped into 7 regions:

- High Income countries (HIGH)
- Low- and middle-income countries of the African Region (AFRO)
- Low- and middle-income countries of the Americas (AMRO)
- Low- and middle-income countries of the Eastern Mediterranean Region (EMRO)
- Low- and middle-income countries of the European Region (EURO)
- Low- and middle-income countries of the South-East Asia Region (SEARO)
- Low- and middle-income countries of the Western Pacific Region (WPRO)

For each region, the available fatality rates of the three economic sectors are shown in Table 2 below.

Table 2: Fatality Occupational Injury Rates per 100,000 employees

Region	Fatality Rates of each economic sector					
	Agriculture		Industry		Service	
	2010	2014	2010	2014	2010	2014
HIGH	7.8	no change	3.8	no change	1.5	no change
AFRO	18.9	no change	21.1	no change	17.7	no change
AMRO	9.3	8.7 ↓	9.5	11.2 ↑	6.0	5.7 ↓
EMRO	13.0	no change	14.9	no change	12.3	no change
EURO	15.7	17.0 ↑	10.3	13.4 ↑	5.5	3.5 ↓
SEARO	24.0	27.5 ↑	9.7	9.9 ↑	5.1	4.4 ↓
WPRO	24.0	27.5 ↑	9.7	9.9 ↑	5.1	4.4 ↓

The previous rates of fatal occupational injuries are used for HIGH, AFRO and EMRO regions because of a lack of data. The percentage of labour force for each economic sector in each country was retrieved from the World Factbook of the Central Intelligence Agency (CIA). These percentages can also be obtained from ILOSTAT Database but they are percentages of the employed instead of the labour force. Together with the labour force, total employment (comprising both paid employment and self-employment) and the respective regions' fatality rates in 2014, the number of fatalities of each country was then computed ^[6].

2.3 Non-fatal Occupational Accidents

As non-fatal (causing at least 4 days of absence) occupational accidents are not usually reported by most countries, they were estimated by using lower and upper limit estimates^[6]. The lower limit of 0.14% was obtained by averaging the proportion of fatal and non-fatal injuries of the European Union (EU) 15 countries except Greece. The upper limit of 0.08% was obtained similarly from Finland, France and Germany. The lower and upper limits used for 2010 were 0.13% and 0.10% respectively. The lower and upper limit estimates of the number of non-fatal injuries of each country in 2014 are then calculated as follows:

$$\text{Estimated number of non fatal injuries (Lower Limit)} = \frac{\text{No. of fatalities} \times 100\%}{0.14}$$

$$\text{Estimated number of non fatal injuries (Upper Limit)} = \frac{\text{No. of fatalities} \times 100\%}{0.08}$$

The estimated non-fatal injury is then finally obtained by taking the mean of the two limits.

2.4 Fatal Work-Related Diseases

The WHO Global Burden of Disease (GBD) is the only data source that covers all the diseases in the world. It categorises all the diseases into seven categories:

- communicable diseases
- malignant neoplasms
- circulatory diseases
- neuropsychiatric conditions
- respiratory diseases
- digestive diseases
- genitourinary diseases.

Since fatal diseases that are work-related are not listed separately in the GBD data, they are estimated by attributable fractions (AF) taken from Nurminen & Karjalainen, 2001.^[5] An attributable fraction can be interpreted as “the fraction of a disease [or injury] which would not have occurred had the exposure factor been non-existent in the population in question.”^[5] The attributable fractions used to calculate the fatal work-related diseases are shown in Table 3. As the attributable fraction for communicable diseases was obtained from industrialized countries in the HIGH region, the attributable fraction was thus reduced for countries that do not belong to this high income group.^[5]

Table 3: Attributable Fractions used in the estimation of fatal work-related diseases

Causes of Death	HIGH Region		OTHER Regions	
	Men	Women	Men	Women
Communicable diseases	4.80	32.50	3.05	20.68
Malignant neoplasms	13.80	2.20	13.80	2.20
Neuropsychiatric conditions	6.60	1.80	6.60	1.80
Circulatory diseases	14.40	6.70	14.40	6.70
Respiratory diseases				
• COPD	18.00	6.00	18.00	6.00
• Asthma	21.00	13.00	21.00	13.00
• Others	1.00	1.00	1.00	1.00
Digestive diseases	2.30	1.50	2.30	1.50
Genitourinary system	3.00	0.40	3.00	0.40

^a Attributable fractions from Nurminen & Karjalainen, 2001

^b Revised attributable fractions

The Collegium Ramazzini released its 19th Statement containing a comprehensive analysis of Chronic Obstructive Pulmonary Disease (COPD) in occupational settings in 2016: A New Approach to the Control of COPD.^[18] As a result, AF for COPD were used to estimate the number of deaths due to respiratory diseases. The AF for COPD was 18% for males and 6% for females.

The estimate of fatal work-related diseases of each WHO region was calculated by selecting the 7 main disease categories with the appropriate AF. The fatal work-related disease of each country was obtained by multiplying the WHO region's estimate with the proportion of its labour force.

3 Results

3.1 Overview

There was an estimated 2.78 million fatalities compared to 2.33 million estimated in 2011. For fatal occupational accidents, there were 380,500 deaths, an increase of 8% in 2014 compared to 2010. Fatal work-related diseases had been a concern since 1998 and were at least five times higher than fatalities due to occupational accidents. In 2015, there were 2.4 million deaths due to fatal work-related diseases, an increase of 0.4 million compared to 2011.

In total, it is estimated that more than 7,500 people die every day; 1,000 from occupational accidents and 6,500 from work-related diseases. The rate of fatal occupational accidents increased slightly. The number of non-fatal occupational accidents was estimated to be 374 million, increasing significantly from 2010. The main reason was that a higher under-reporting estimate was used compared to the previous estimates.

Table 4: Global Trend of Occupational Accidents and Fatal Work-Related Diseases (1998-2015)

Year	Fatal Occupational Accidents		Non-fatal Occupational Accidents At least 4 days absence		Fatal Work-Related Disease
	Number	Rate ^a	Number	Rate ^a	
1998	345,436	16.4	263,621,966	12,534	
2000					2,028,003
2001	351,203	15.2	268,023,272	12,218	
2002					1,945,115
2003	357,948	13.8	336,532,471	12,966	
2008	320,580	10.7	317,421,473	10,612	2,022,570
2010	352,769	11.0	313,206,348	9,786	
2011					1,976,021
2014	380,500	11.3	373,986,418	11,096	
2015					2,403,965

^a Number of occupational accidents per 100,000 persons in the labour force

3.2 Occupational Accidents

3.2.1 By WHO Region

Table 5: Breakdown of 2014 Occupational Accidents by WHO Regions

Region	Labour force	Total employment	Occupational injuries reported to ILO in 2014		Global estimates of occupational accidents 2014			
			Fatal	Non-fatal	Fatal	Non-fatal (at least four days absence)		
						Lower limit (0.14)	Upper limit (0.08)	Average
HIGH	521,662,897	476,238,830	4,064	1,991,628	10,757	7,683,291	13,193,091	10,845,213
AFRO	350,749,965	345,598,985	110	8,742	65,145	46,532,038	81,431,066	63,981,552
AMRO	297,081,063	269,154,947	1,916	966,221	19,388	13,848,697	24,235,220	19,041,958
EMRO	195,451,073	160,588,073	210	16,692	21,113	15,081,021	26,391,787	20,736,404
EURO	227,406,974	204,171,210	2,598	98,904	14,159	10,113,353	17,698,368	13,905,861
SEARO	824,496,607	763,156,900	77	6,710	124,404	88,859,942	155,504,898	122,182,420
WPRO	953,638,990	909,608,518	222	44,638	125,535	89,667,644	156,918,377	123,293,011
TOTAL	3,370,487,570	3,128,517,463	9,197	3,133,535	380,500	271,785,986	475,372,807	373,986,418

Similar to 2010, most of the occupational accidents in 2014 had occurred in the SEARO and WPRO regions as shown in Table 5. About two-thirds of the occupational accidents fell almost equally under these two regions. Compared to the 2010 figures, there was a rise in the number of occupational accidents for all the WHO regions except for the HIGH and EURO regions.

3.2.2 By Geographical Region

The number of fatal and non-fatal occupational accidents of the 5 geographical regions are presented in Table 6. Asia had the highest number of fatalities among the 5 regions and constituted more than 70% globally. The Asian fatal occupational rate was 12.7 per 100,000 persons in the labour force which was lower than Africa which had the highest fatality rate of 16.6 per 100,000 persons in the labour force. Europe had the lowest fatality rate among the 5 regions, with a rate of 3.61.

Table 6: Breakdown of 2014 Occupational Accidents by Geographical Regions


Region	Labour force	Total employment	Occupational injuries reported to ILO 2014		Global estimates of occupational accidents 2014				
			Fatal	Non-fatal	Fatal	* Fatality Rate	Non-fatal (at least four days absence)		
							Lower limit (0.14)	Upper limit (0.08)	Average
Africa	413,329,046	397,013,885	320	25,434	71,882	17.39	51,343,960	89,851,931	70,597,946
America	479,990,599	433,527,137	1,916	966,221	24,579	5.12	17,556,317	30,672,033	24,114,175
Asia	2,093,134,505	1,953,718,973	2,694	121,256	271,949	12.99	194,249,063	339,932,172	267,097,755
Europe	364,546,703	326,139,450	4,079	1,921,644	11,017	3.02	7,869,606	13,600,463	11,134,918
Oceania	19,486,718	18,118,018	188	98,980	1,074	5.51	767,040	1,316,209	1,041,625
Total	3,370,487,570	3,128,517,463	9,197	3,133,535	380,500	11.29	271,785,986	475,372,807	373,986,418

* Number of occupational accidents per 100,000 persons in the labour force

3.3 Work-Related Diseases

3.3.1 Work-related Mortality by Cause

Figure 1: Breakdown of the Estimated Fatal Work-Related Mortality by Cause in 2015



In Figure 1, the main causes of death from all work-related diseases were circulatory diseases (31%), work-related cancers (26%), respiratory diseases (17%) and occupational injuries (14%). They formed about 90% of all fatal work-related deaths. Table 6 showed the work-related diseases by WHO regions. Communicable diseases were still a common problem in developing countries. The current burden of 9% had decreased slightly from 10% in 2011 probably due to better estimates since the estimate in 2008 was 25% which was conservative.

Figure 2: Fatal Work-Related Mortality by WHO Regions and Cause in 2015





Figure 3: Comparison of Fatal Work-Related Mortality by WHO Regions between 2011 and 2015


In Figure 3, communicable diseases constituted slightly more than 30% in AFRO in 2015 among the work-related deaths but less than 5% in HIGH. The proportion of the cause of fatal work-related diseases had remained almost the same except for an increased proportion of respiratory diseases occurring in all the regions. Previous respiratory disease estimates were clearly underestimates as they were lower than respiratory deaths attributed to hazardous substances. On the other hand, circulatory diseases had increased in all regions except for HIGH and EURO.

Although the number of deaths due to neuropsychiatric conditions was low (2%), it had increased slightly in 2015. It could be assumed that the number of deaths due to neuropsychiatric conditions will increase in the future. These diseases, together with musculoskeletal disorders, are usually non-fatal and are relatively more prevalent in developed countries.

The proportion of work-related cancer (malignant neoplasm) dominated the fatal work-related diseases in high income countries. This is a continuously growing problem that needs better preventive measures. Practically, all such cancers can be avoided by eliminating the exposures to carcinogenic agents and minimizing exposures at related jobs, settings and procedures.

Table 7: Breakdown of the Estimated Fatal Work-Related Mortality by WHO Regions in 2015

	HIGH	AFRO	AMRO	EMRO	EURO	SEARO	WPRO	TOTAL
Communicable disease	10,435	84,948	8,152	19,396	4,373	87,511	15,168	229,983
Malignant neoplasms	225,939	37,198	50,038	29,036	56,277	110,662	233,085	742,235
Neuropsychiatric conditions	25,512	3,927	3,390	2,396	1,101	7,280	4,510	48,116
Circulatory diseases	103,863	60,151	48,580	50,597	129,992	246,885	223,105	863,173
Respiratory diseases	51,363	21,419	18,834	27,123	13,714	215,118	128,018	475,589
Digestive diseases	3,132	4,505	2,535	1,546	2,910	8,631	2,655	25,914
Genitourinary diseases	3,840	1,750	1,835	1,167	729	7,214	2,420	18,955
Occupational injuries	10,757	65,145	19,388	21,113	14,159	124,404	125,535	380,500
TOTAL	434,840	279,042	152,754	152,375	223,253	807,705	734,497	2,784,465

3.3.2 Number of Work-related Mortality by United Nations Geographical Regions


Table 8 showed the estimated work-related mortality by the United Nations geographical regions. Asia had the highest number of work-related mortality among the five regions. About two-thirds of the work-related mortalities came from Asia (Figure 4). In fact, Asia had in absolute numbers, the highest number of work-related deaths in each disease category, including occupational injuries.

Table 8: Breakdown of Work-Related Mortality by United Nations Geographical Regions in 2015

	AFRICA	AMERICA	ASIA	EUROPE	OCEANIA	* TOTAL
Communicable Diseases	91,158	11,754	119,397	7,075	374	229,758
Malignant neoplasms	46,494	128,020	430,472	126,127	7,645	738,759
Neuropsychiatric conditions	4,694	12,195	19,856	10,540	762	48,048
Circulatory diseases	76,350	84,428	563,687	131,335	4,052	859,853
Respiratory diseases	30,103	36,562	375,611	29,322	2,086	473,683
Digestive diseases	5,000	3,616	13,906	3,249	103	25,874
Genitourinary diseases	2,123	3,161	11,529	1,983	123	18,919
Occupational injuries	71,905	23,101	268,641	14,094	895	378,634
Total	327,827	302,837	1,803,098	323,726	16,040	2,773,528

* The numbers did not tally with Table 6 due to rounding off errors.

Figure 4: Distribution of Work-related Mortality by UN Geographical Regions



3.3.3 Deaths attributed to hazardous substances including dusts, vapours and fumes

Table 9: Estimated deaths attributed to hazardous substances (incl. dusts, vapours and fumes)

	No. of deaths		Estimated % attributed to hazardous substances		No. of deaths attributed to hazardous substances		
	Men	Women	Men	Women	Men	Women	Total
Cancers	4,779,542	3,754,270	-	-	343,646	71,976	415,622
Mouth and oropharynx cancers	242,334	96,859	1.0	0.5	2,423	484	2,908
Oesophagus cancer	321,263	136,534	3.3	1.1	10,602	1,502	12,104
Stomach cancer	499,852	292,749	3.0	0.3	14,996	878	15,874
Colon and rectum cancers	396,370	349,958	1.0	0.5	3,964	1,750	5,713
Liver cancer	559,329	255,598	0.2	0.1	1,119	256	1,374
Pancreas cancer	169,105	161,678	0.02	0.01	34	16	50
Trachea, bronchus, lung cancers	1,113,475	520,462	21.1	5.3	234,943	27,585	262,528
Melanoma and other skin cancers	54,769	40,412	10.0	2.0	5,477	808	6,285
Breast cancer	0	558,514	-	4.6	0	25,692	25,692
Cervix uteri cancer	0	306,319	-	0.7	0	2,144	2,144
Ovary cancer	0	161,181	-	0.5	0	806	806
Prostate cancer	328,131	0	1.0	-	3,281	0	3,281
Bladder cancer	143,275	47,731	7.1	1.9	10,173	907	11,079
Leukaemia	136,879	109,581	0.9	0.5	1,232	548	1,780
Other malignant neoplasms	814,759	716,693	6.8	1.2	55,404	8,600	64,004
Neuropsychiatric conditions	82,630	75,083	1.0	1.0	826	751	1,577
Cardiovascular diseases	4,366,727	3,497,981	1.0	1.0	43,667	34,980	78,647
Respiratory Diseases	2,259,742	1,867,406	-	-	362,910	112,679	475,589
Chronic obstructive pulmonary disease	1,740,463	1,443,292	18.0	6.0	313,283	86,598	399,881
Asthma	222,168	182,004	21.0	13.0	46,655	23,660	70,316
Other respiratory diseases	297,112	242,110	1.0	1.0	2,971	2,421	5,392
Genitourinary diseases	561,351	526,893	1.0	1.0	5,614	5,269	10,882
Total					756,663	225,654	982,317

In 2015, nearly one million workers died at work due to hazardous substances (Table 9), an increase of more than 90,000 workers when compared to 2011.

4 Discussion

The methodology used to calculate the estimates was similar to the previous global estimates. The 2014 total employment was collected from ILOSTAT. Total employment is defined to be the number of persons employed including paid employment and self-employment. If total employment numbers were not available, the 2010 numbers were used instead. If neither were available, labour force numbers were used. Since the affected countries are usually small, there would be minimal impact on the global figures.

Labour Force data was collected from ILOSTAT. For ILO countries where ILOSTAT was not available, it was obtained from the earlier LABORSTA data also from ILO. For non-ILO countries, the figures were obtained from CIA World Factbook. For countries where such data was not available, 2010 estimates were used instead.

The percentage of the total employment by sector of economic activity was collected from CIA World Factbook. For countries where such data was not available, previous estimates were used instead. Once the fatal occupational injury rate of each economic sector in Section 2.1 was known, fatal occupational injuries could then be calculated. As new fatal occupational injury rates were only available for AMRO, EURO and WPRO from ILOSTAT, 2010 estimates were then used for HIGH, AFRO and EMRO. For AMRO, Chile (2011-2013) and Colombia (2015) were used to formulate the fatal occupational injury rates for agriculture, industry and service. For EURO, Russia (2015) and Ukraine (2012) were used. For WPRO, Macau (2015) and Malaysia (year 2015) were used. Though these representative countries were not used during the previous estimates, the calculated rates did not differ too much from the previous rates as shown in Table 1. For SEARO, the rates for WPRO were used.

There was an increase in employment for the seven WHO regions. Though the total employment in 2014 for all countries increased by 4% compared to 2010, the number of estimated fatal occupational accidents increased by about 8% to 380,500. However, the number of fatal occupational accidents decreased in two of the WHO regions despite an increase in the total employment. There was an estimated 6% decrease in occupational accidents for HIGH Region despite a 5% increase in total employment. For EURO, occupational accidents decreased by 3% with a 2% increase in employment. However for WPRO and SEARO, though employment increased by 2-3%, the number of fatal occupational accidents was estimated to increase by 8-9%. As such, the estimated fatal and non-fatal occupational injury rates in 2014 had increased and was higher than the global crisis year in 2008 (Table 3).

Chronic Obstructive Pulmonary Disease (COPD) is a growing disease world-wide. It has become an important work-related problem but is often not well-recognized and is thus under-diagnosed. It is caused by exposures to a multitude of vapours, gases, dusts and fumes which are collectively known as VGDF (vapours, gases, dusts and fumes). Many VGDF exposures pose serious risks in workplaces and are often under-regulated. Besides VGDF, COPD is known to be associated with specific occupational exposure agents including coal mine dust, asbestos, silica, welding and cutting gases and fumes, cement dust, diesel exhausts, spray painting, organic solvents and possibly man-made mineral fibres¹⁸.

The American Thoracic Society estimated that occupational exposures were responsible for 15-20% of COPD in 2004 and this had since been supported by many publications that occupational exposures were an important contribution to COPD. In light of this, the previous respiratory death numbers were clearly underestimates as also evidenced in Table 9 for hazardous substances. For example, in the previous estimate, the estimated deaths due to COPD was 374,128 which was more than twice the number of work-related diseases due to respiratory diseases at 162,260. For the computation of the respiratory estimates, AF values were used to calculate the number of deaths due to COPD. They were 18% for males and 6% for females. Deaths due to COPD constituted more than 80% of the respiratory estimates.

Among the more than 400,000 increase in fatal work-related diseases in 2015, more than 70% were due to respiratory diseases. Estimates of fatal work-related diseases due to respiratory diseases is now the third largest after circulatory diseases and malignant neoplasms. A new paradigm has to be adopted to recognise that occupational exposures to VGDF cause COPD and preventive measures should be taken to reduce such exposures to workers.

References

- ¹ Takala, J. "ILO Introductory Report: Decent Work - Safework". ILO Introductory Report, XVII World Congress on Safety and Health at Work, Orlando Florida, ISBN 92-2-117750-5 http://ohsa.org.mt/Portals/0/docs/intrep_05.pdf (accessed on 15 October 2015)
- ² ILO 2014. Safety and Health at Work: a vision for sustainable prevention. XX World Congress on Safety and Health at Work, 2014, Global Forum for Prevention, Frankfurt. International Labour Office, Geneva, 41 p. <https://goo.gl/RxXHhU> (accessed on 16 Nov 2016)
- ³ National Bureau of Statistics of China. Statistical Communiqué of the People's Republic of China on the 2013 National Economic and Social Development, Chapter XII. Resources, Environment and Work Safety http://www.stats.gov.cn/english/PressRelease/201402/t20140224_515103.html (accessed 13 Dec.2015)
- ⁴ Nurminen M & Karjalainen A (2001). Epidemiologic estimate of the proportion of fatalities related to occupational factors in Finland. *Scandinavian Journal of Work, Environment & Health* 27, 161–213. http://www.sjweh.fi/show_abstract.php?abstract_id=605. (accessed on 26 March 2014)
- ⁵ Hämäläinen P. Global Estimates of Occupational Accidents and Fatal Work-Related Diseases. Doctoral dissertation, Publication 917, Tampere University of Technology, Finland, 2010. web page: <http://dspace.cc.tut.fi/dpub/bitstream/handle/123456789/6818/hamalainen.pdf?sequence=1>. (accessed on 7 Dec 2016)
- ⁶ Hämäläinen, P., Takala, J., Saarela, K.L. 2006. Global estimates of occupational accidents. *Safety Science* 44, 137–156, web page: http://www.bvsde.paho.org/bvsacd/cd63/global_estimates.pdf
- ⁷ Ana M García, Rafael Gadena Merino, Vicente López Martínez: Estimación de la mortalidad atribuible a enfermedades laborales en España, 2004. *Rev Esp Salud Pública* 2007; 81 :261-270, N.º3 Mayo-Junio 2007. Accessible from: http://scielo.isciii.es/scielo.php?pid=S1135-57272007000300003&script=sci_arttext (accessed on 15 September 2014)
- ⁸ Andrea 't Mannelje, Neil Pearce: Quantitative estimates of work-related deaths, diseases and injury in New Zealand. *Scand J Work Environ Health* 2005;31(4):266-276; doi:10.5271/sjweh.882. Accessible from www.sjweh.fi/download.php?abstract_id=882&file_nro=1 (accessed on 15 September 2014)
- ⁹ Steenland K, Rollins School Public Health, Emory University, Atlanta, Ga. USA. 2004 <http://www.occupationalcancer.ca/wp-content/uploads/2011/03/Steenland.pdf> (accessed on 14 September 2014)
- ¹⁰ Lesley Rushton, Sally J Hutchings, Lea Fortunato, Charlotte Young, Gareth S Evans, Terry Brown, Ruth Bevan, Rebecca Slack, Phillip Holmes, Sanjeev Bagga, John W Cherrie and Martie Van Tongeren. Occupational cancer burden in Great Britain *Br J Cancer* 107: S3-S7; doi:10.1038/bjc.2012.112 <http://www.nature.com/bjc/journal/v107/n1s/index.html> (Accessed on 16 May 2013)

¹¹ Driscoll, T., J. Takala, K. Steenland, C. Corvalan, and M. Fingerhut. Review of estimates of the global burden of injury and illness due to occupational exposures. *Am. J. Ind. Med.* 48:491–502 (2005).

¹² Takala J, Hämmäläinen P, Saarela KL, Loke YY, Manickam K, Tan WJ, Heng P, Tjong C, Lim GK, Lim S, Gan SL: Global Estimates of the Burden of Injury and Illness at Work in 2012. *JOEH* 11: 326-337, May/2014, Taylor & Francis, open access, <http://www.tandfonline.com/doi/pdf/10.1080/15459624.2013.863131>

¹³ Nenonen N, Hämmäläinen P, Takala J, Saarela KL, Yong E, Lim SL, Lim GK, Manickam K: Global Estimates of Occupational Accidents and Fatal Work-related Diseases in 2014, based on 2010 and 2011 Data from ILO and WHO, Report to the ILO, Tampere, Singapore, Geneva 2014, DOI: 10.13140/2.1.2864.0647, web page: <http://goo.gl/aaJY9r>


¹⁴ Stephen S Lim, Theo Vos, Abraham D Flaxman, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters I 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; 380: 2224-60, Institute of Health Metrics and Evaluation web site: <http://vizhub.healthdata.org/gbd-cause-patterns/>

¹⁵ Hämmäläinen P. Global Estimates of Occupational Accidents and Fatal Work-Related Diseases. Doctoral dissertation, Publication 917, Tampere University of Technology, Finland, 2010. Accessed on 11 September 2014 and available at <http://dspace.cc.tut.fi/dpub/bitstream/handle/123456789/6818/hamalainen.pdf?sequence=1>

¹⁶ GBD 2015 Mortality and Causes of Death Collaborators. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016;388:1459-544. See details from <http://vizhub.healthdata.org/gbd-compare/>

¹⁷ Takala J, Young E. Mortality estimates of occupational injuries and ill health in Singapore. WSH Institute, Public Health and Occupational Medicine Conference, 2014 Singapore. <https://goo.gl/RqvCKp> (accessed on 17 Nov 2016)

¹⁸ Collegium Ramazzini 19th Statement: A New Approach to the Control of Chronic Obstructive Pulmonary Disease (COPD). 19 July 2016, <http://www.collegiumramazzini.org/news1.asp?id=144> and [http://www.collegiumramazzini.org/download/19_NineteenthCRStatement\(2016\).pdf](http://www.collegiumramazzini.org/download/19_NineteenthCRStatement(2016).pdf) (accessed 7 Dec 2016)



Workplace Safety and Health Institute
1500 Bendemeer Road, #04-01
Ministry of Manpower Services Centre
Singapore 339946

www.wsh-institute.sg

ISBN: 9789811148446